

Erreger	Testverfahren
<i>B. henselae</i>	Indirekter Immunfluoreszenztest
<i>B. pertussis</i>	IgG-EIA IgA-EIA
Borrelien-Serologie <i>B. burgdorferi sensu lato</i>	Serum-IgG-recom-EIA Serum-IgM-recom-EIA Liquor-IgG-recom-EIA Liquor-IgM-recom-EIA Serum und Liquor-IgG-line-Blot Serum und Liquor-IgM-line-Blot IgG-Serum-Liquor-Quotient
<i>Brucella spp.</i>	IgM und IgG-EIA
<i>Campylobacter spp.</i>	IgM und IgG rek. Line Blot
<i>C. pneumoniae</i>	IgG-EIA IgM-EIA
<i>C. trachomatis</i>	IgG-EIA
<i>C. burnettii</i>	IgM und IgG-EIA
<i>C. tetani</i>	s. Impfserologie
<i>C. diphtheriae</i>	s. Impfserologie
<i>H. influenzae B</i>	s. Impfserologie
<i>L. interrogans</i>	IgM und IgG-EIA
<i>M. pneumoniae</i>	IgG/IgM/IgA-EIA
<i>M. tuberculosis</i>	QuantiFERON® -TB Gold Plus
<i>Pseudomonas</i>	IgG-EIA
<i>S. aureus</i>	Anti-Staphylolysin (Agglutinationstest)
<i>Streptococcus pneumoniae</i>	IgG-AK
Lueserologie <i>T. pallidum</i>	CLIA / TPHA Serum TPHA (Indirekter Hämagglutinationstest) Liquor FTA-Abs IgM/ IgG (IFT) VDRL (Agglutinationstest) Serum VDRL (Agglutinationstest) Liquor Serum/Liquor -IgG-Line-Blot Serum/ Liquor-IgM-Line-Blot Serum-Liquor Quotient IgM-EIA
<i>Yersinia spp.</i>	IgG-EIA IgA-, IgM- und IgG-EIA IgG/IgM/ IgA recom Blot

Pilze	Testverfahren
<i>Aspergillus spp.</i> (1→3)-β-D-Glucan <i>C. albicans</i> <i>C. neoformans</i> <i>Histoplasma spp.</i> <i>Blastomyces spp.</i> <i>Coccidiomyces spp.</i>	Ag - EIA
	Immunpräzipitation Ouchterlony
	Ag-EIA
	Agglutinationstest
	Immunpräzipitation Ouchterlony
	Immunpräzipitation Ouchterlony
	Immunpräzipitation Ouchterlony

Wurminfektionen	Testverfahren
<i>Echinococcus spp.</i> <i>Toxocara canis</i>	<i>E. granulosus</i> -IHA
	<i>E. spec.</i> -EIA
	<i>E. multilocularis</i> -EIA
	<i>Echinococcus spp.</i>
	Immunoblot
	Immunoblot

Protozoen	Testverfahren
<i>T. gondii</i>	IgG-EIA IgM-EIA IgG-Avidität